

Skyrmions in Spinor Bose-Einstein Condensates

H.T.C. Stoof

*Institute for Theoretical Physics, University of Utrecht, Princetonplein 5,
3584 CC Utrecht, The Netherlands*

We show that spinor Bose-Einstein condensates not only have line-like vortex excitations, but in general also allow for point-like topological excitations, i.e., skyrmions. We discuss the static and dynamic properties of these skyrmions for spin-1/2 and ferromagnetic spin-1 Bose gases.

PACS number(s): 03.75.Fi, 67.40.-w, 32.80.Pj

Introduction. — An understanding of quantum magnetism is important for a large number of phenomena in physics. Three well-known examples are high-temperature superconductivity, quantum phase transitions and the quantum Hall effect. Moreover, it appears that magnetic properties will also be very important in another area, namely Bose-Einstein condensation in trapped atomic gases. The latter has come about because of two independent experimental developments. The first development is the realization of an optical trap for atoms, whose operation no longer requires the gas to be doubly spin-polarized [1,2]. The second development is the creation of a two-component Bose condensate [3], which by means of rf-fields can be manipulated so as to make the two components essentially equivalent [4]. As a result the behavior of both spin-1 and spin-1/2 Bose gases can now be experimentally explored in detail.

Theoretically, the ground-state structure of these gases has recently been worked out by a number of authors [5–8] and also the first studies of the line-like vortex excitations have appeared [5,9]. However, an immediate question that comes to mind is whether the spin degrees of freedom allow for other topological excitations that do not have an analogy in the case of a single component Bose condensate. It is the main purpose of this Letter to show that the answer to this question is in general affirmative. In particular, we show that spinor Bose-Einstein condensates have so-called skyrmion excitations, which are topological nontrivial point-like spin textures [10]. Having done so, we then turn to the investigation of the precise texture and the dynamics of these skyrmions.

Topological considerations. — To find all possible topological excitations of a spinor condensate, we need to know the full symmetry of the macroscopic wavefunction $\Psi(\mathbf{x}) \equiv \sqrt{n(\mathbf{x})}\zeta(\mathbf{x})$, where $n(\mathbf{x})$ is the total density of the gas, $\zeta(\mathbf{x})$ is a normalized spinor that determines the average local spin by means of $\langle \mathbf{S} \rangle(\mathbf{x}) = \zeta_a^*(\mathbf{x}) \mathbf{S}^{ab} \zeta_b(\mathbf{x})$, and $\mathbf{S}$ are the usual spin matrices obeying the commutation relations $[S_\alpha, S_\beta] = i\epsilon_{\alpha\beta\gamma} S_\gamma$. Note that here, and in the following, summation over repeated indices is always implicitly implied. From the work of Ho [5] we know that in the case of spin-1 bosons we have to consider two possibilities, since the effective interaction between two spins can be either antiferromagnetic or ferromagnetic.

In the antiferromagnetic case the ground-state energy is minimized for $\langle \mathbf{S} \rangle(\mathbf{x}) = \mathbf{0}$, which implies that the parameter space for the spinor $\zeta(\mathbf{x})$ is only $U(1) \times S^2$ because we are free to choose both its overall phase and the spin quantization axis. In the ferromagnetic case the energy is minimized for $|\langle \mathbf{S} \rangle(\mathbf{x})| = 1$ and the parameter space corresponds to the full rotation group $SO(3)$. Using the same arguments, we find that for spin-1/2 bosons the order parameter space of the ground state is always equivalent to $SU(2)$ [11].

What do these results tell us about the possible topological excitations? For line-like defects or vortices, we can assume $\zeta(\mathbf{x})$ to be independent of one direction and the spinor represents a mapping from a two-dimensional plane into the order parameter space. If the vortex is singular this will be visible on the boundary of the two-dimensional plane and we need to investigate the properties of a continuous mapping from a circle S^1 into the order parameter space G , i.e., from the first homotopy group $\pi_1(G)$. Since $\pi_1(SU(2)) = \pi_1(SO(3)) = Z_2$ and $\pi_1(U(1) \times S^2) = Z$, we conclude that a spin-1/2 and a ferromagnetic spin-1 condensate can have only vortices with a winding number equal to 1, whereas an antiferromagnetic spin-1 condensate can have vortices with winding numbers that are an arbitrary integer. Physically, this means that by traversing the boundary of the plane, the spinor can wind around the order parameter at most once or an arbitrary number of times, respectively. This conclusion is identical to the one obtained previously by Ho [5].

If the vortex is nonsingular, however, the spinor $\zeta(\mathbf{x})$ will be identical everywhere on the boundary of the two-dimensional plane and it effectively represents a mapping from the surface of a three-dimensional sphere S^2 into the order parameter space. We then need to consider the second homotopy group $\pi_2(G)$. For this we have that $\pi_2(SU(2)) = \pi_2(SO(3)) = 0$ and $\pi_2(U(1) \times S^2) = Z$. Hence nonsingular or coreless vortices are only possible for a spin-1 condensate with antiferromagnetic interactions. It therefore appears that the nonsingular spin texture discussed in Ref. [5], is topologically unstable and can be continuously deformed into the ground state by ‘local surgery’ [12].

We are now in a position to discuss point-like de-

fects. Since the boundary of a three-dimensional gas is also the surface of a three-dimensional sphere, singular point-like defects are also determined by the second homotopy group $\pi_2(G)$. Such topological excitations thus only exist in the case of a spin-1 Bose gas with antiferromagnetic interactions. We call these excitations singular skyrmions, although in view of the work of 't Hooft and Polyakov it would be justified to call them monopoles [13]. For nonsingular point-like defects the spinor $\zeta(\mathbf{x})$ will again be identical on the boundary of the three-dimensional gas. As a result, the configurations space is compactified to the surface of a four-dimensional sphere S^3 and we need to determine the third homotopy group $\pi_3(G)$. For this we find $\pi_3(SU(2)) = \pi_3(SO(3)) = \pi_3(U(1) \times S^2) = Z$. Hence nonsingular skyrmion excitations exist in all three cases.

Skyrmion texture. — In view of the fact that magnetic gradients are needed to stabilize the spinor condensate in the case of a spin-1 Bose gas with antiferromagnetic interactions [8], we from now on consider only spin-1/2 and ferromagnetic spin-1 Bose gases, which have only nonsingular skyrmions as we have seen. Due to the absence of a core, the fluctuations in the density $\delta n(\mathbf{x}) = n(\mathbf{x}) - \langle n(\mathbf{x}) \rangle$ will be small compared to the average density $\langle n(\mathbf{x}) \rangle$ and the energy of the skyrmion can be determined by

$$E[\zeta] = \frac{1}{2} \int d\mathbf{x} \int d\mathbf{x}' \delta n(\mathbf{x}) \chi^{-1}(\mathbf{x}, \mathbf{x}') \delta n(\mathbf{x}') + \int d\mathbf{x} n(\mathbf{x}) \left(\frac{\hbar^2}{2m} |\nabla \zeta(\mathbf{x})|^2 - \mu \mathbf{B} \cdot \langle \mathbf{S} \rangle(\mathbf{x}) \right). \quad (1)$$

Here, $\chi(\mathbf{x}, \mathbf{x}')$ is the static density-density response function defined by

$$\frac{\hbar^2}{4m} \left(-\nabla \cdot \left(\frac{1}{\langle n(\mathbf{x}) \rangle} \nabla \right) + 16\pi a \right) \chi(\mathbf{x}, \mathbf{x}') = \delta(\mathbf{x} - \mathbf{x}'), \quad (2)$$

a is the appropriate scattering length, $\mathbf{B}$ is either a fictitious magnetic field, caused by resonant rf-fields, or a real homogeneous magnetic field and μ is the corresponding magnetic moment of the atoms in the trap. Moreover, the spinor $\zeta(\mathbf{x})$ can now be conveniently parametrized as

$$\zeta(\mathbf{x}) = \exp \left\{ -\frac{i}{S} \Omega^\alpha(\mathbf{x}) S_\alpha \right\} \zeta^Z, \quad (3)$$

where S denotes the spin of the atoms and ζ^Z is a constant spinor that minimizes the Zeeman energy, i.e., it obeys $\zeta_a^Z = \delta_{aS}$ if we use the direction of the magnetic field as the quantization axis.

This last equation explicitly shows that the topology of the order parameter space is a sphere of radius π with opposite points on the surface identified. If we now assume that a maximally symmetric shape of the skyrmion spin texture is allowed, we can take $\Omega(\mathbf{x}) = \mathbf{x}\omega(x)/x \equiv$

$\hat{\mathbf{x}}\omega(x)$, with $\omega(x)$ a monotonically decreasing function obeying $\omega(0) = 2\pi$ and $\lim_{x \rightarrow \infty} \omega(x) = 0$. For this ansatz we see that by traversing the whole configuration space, we exactly cover the order parameter space once. The above spin texture therefore corresponds to a skyrmion with a topological winding number

$$W = \frac{3}{4\pi^4} \int d\Omega = \frac{1}{8\pi^4} \int d\mathbf{x} \epsilon^{ijk} \epsilon_{\alpha\beta\gamma} \partial_i \Omega^\alpha \partial_j \Omega^\beta \partial_k \Omega^\gamma \quad (4)$$

equal to 1 and located at the center of the trap.

To obtain the precise spin texture of the skyrmion we have to find the function $\omega(x)$ that minimizes the energy. In general this leads, due to the nonlocality of the density-density response function, to a complicated nonlinear integro-differential equation, which can only be solved numerically. However, we can gain a lot of physical insight in the size and stability of the skyrmion by the following analysis. Considering first the ideal case of zero magnetic field $\mathbf{B}$ and solving for the density fluctuations induced by the spin texture, we have to minimize the gradient energy

$$E^{\text{grad}}[\zeta] = \int d\mathbf{x} \langle n(\mathbf{x}) \rangle \frac{\hbar^2}{2m} |\nabla \zeta(\mathbf{x})|^2 - \frac{\hbar^4}{8m^2} \int d\mathbf{x} \int d\mathbf{x}' |\nabla \zeta(\mathbf{x})|^2 \chi(\mathbf{x}, \mathbf{x}') |\nabla \zeta(\mathbf{x}')|^2. \quad (5)$$

For a skyrmion of size λ in the center of the trap this energy can be estimated to be given by

$$E^{\text{grad}}(\lambda) \simeq \langle n(\mathbf{0}) \rangle \frac{2\pi\hbar^2}{m} \left(\lambda - \frac{3\lambda}{(\lambda/\xi)^2 + 1} \right), \quad (6)$$

with $\xi = (16\pi|a|\langle n(\mathbf{0}) \rangle)^{-1/2}$ the typical coherence length of the condensate. It has a minimum when λ/ξ is equal to $((\sqrt{33} - 5)/2)^{1/2} \simeq 0.61$. Therefore, our maximally symmetric ansatz is indeed justified, even for an anisotropic trap, as long as the coherence length is much smaller than the size of the condensate, i.e., we are in the so-called Thomas-Fermi limit.

For this result we have assumed the scattering length a to be positive. In that case the skyrmion is thus stabilized by the fact that gradients in the spin texture on the one hand lead to an increase in the average kinetic energy of the spinor condensate, but on the other hand also lead to a reduction of the density, which for repulsive interactions results in a decrease in the average interaction energy. For a negative scattering length, gradients in the spin texture lead to an enhancement of the density, but this, due to the attractive interactions, also reduces the average interaction energy. Therefore, we expect skyrmions also to be stabilized in that case. Indeed, using the same estimates as before, we find that $E^{\text{grad}}(\lambda)$ is now minimized for λ/ξ equal to $((\sqrt{33} + 5)/2)^{1/2} \simeq 2.3$. However, due to the intrinsic instability of a condensate with attractive interactions [14], this implies that the size

of the skyrmion is essentially equal to the size of the condensate itself and that its spin texture will be strongly affected by finite-size effects. In particular, it in general becomes anisotropic.

Note finally, that the above mechanism for the stability of the skyrmion excitations is quite different from that in the quantum Hall effect near filling fraction $\nu = 1$. There skyrmions are electrically charged and their size is determined by a competition between the Coulomb and Zeeman interactions [15]. In the case of a spinor Bose-Einstein condensate, the Zeeman interaction plays a much less important role and for small magnetic fields $B \ll 8\pi|a|\hbar^2\langle n(\mathbf{0})\rangle/m\mu$ only directs the spins at large distances of the skyrmion. Larger magnetic fields will tend to decrease the size of the skyrmion so as to reduce the cost in Zeeman energy.

Skyrmion dynamics. — The most important dynamical variable of the skyrmion arises from the fact that the Euler-Lagrange equations for the skyrmion spin texture is invariant under a space independent rotation of the average local spin $\langle \mathbf{S} \rangle(\mathbf{x})$ around the magnetic field direction $\hat{\mathbf{B}}$ [16]. Mathematically, this means that if $\zeta^{\text{sk}}(\mathbf{x})$ describes a skyrmion, then $\exp\{-i\vartheta \hat{\mathbf{B}} \cdot \mathbf{S}\} \zeta^{\text{sk}}(\mathbf{x})$ describes also a skyrmion with the same winding number and energy. The dynamics of the variable $\vartheta(t)$ associated with this symmetry is determined by the full action for the spin texture $S[\zeta] = \int dt (T[\zeta] - E[\zeta])$, which contains the time-derivative term

$$T[\zeta] = \int d\mathbf{x} n(\mathbf{x}, t) \zeta^*(\mathbf{x}, t) i\hbar \frac{\partial}{\partial t} \zeta(\mathbf{x}, t). \quad (7)$$

Hence, substituting $\zeta(\mathbf{x}, t) = \exp\{-i\vartheta(t) \hat{\mathbf{B}} \cdot \mathbf{S}\} \zeta^{\text{sk}}(\mathbf{x})$ and making use of the conservation of total particle number N to introduce the change of the average local spin projection on the magnetic field $\langle \Delta S_{||} \rangle(\mathbf{x}) = \hat{\mathbf{B}} \cdot \langle \mathbf{S} \rangle(\mathbf{x}) - S$ induced by the skyrmion, we obtain apart from an unimportant boundary term that

$$T[\zeta] = \hbar \frac{\partial \vartheta(t)}{\partial t} \int d\mathbf{x} n(\mathbf{x}, t) \langle \Delta S_{||} \rangle(\mathbf{x}). \quad (8)$$

Taking now again the density fluctuations into account, we finally find that the dynamics of the rotation angle $\vartheta(t)$ is determined by the action

$$S[\vartheta] = \int dt \left\{ \frac{\partial \vartheta(t)}{\partial t} \hbar \langle \Delta S_{||}^{\text{tot}} \rangle + \frac{1}{2} I \left(\frac{\partial \vartheta(t)}{\partial t} \right)^2 \right\}, \quad (9)$$

where $\langle \Delta S_{||}^{\text{tot}} \rangle$ is the change of the total spin along the magnetic field direction and the ‘moment of inertia’ of the skyrmion equals

$$I = \hbar^2 \int d\mathbf{x} \int d\mathbf{x}' \langle \Delta S_{||} \rangle(\mathbf{x}) \chi(\mathbf{x}, \mathbf{x}') \langle \Delta S_{||} \rangle(\mathbf{x}'). \quad (10)$$

The importance of this result is twofold. First, from the action in Eq. (9) we see that at the quantum level

the hamiltonian for the dynamics of the wave function $\Psi(\vartheta, t)$ becomes

$$H = \frac{1}{2I} \left(\hbar \frac{\partial}{\partial \vartheta} - \hbar \langle \Delta S_{||}^{\text{tot}} \rangle \right)^2. \quad (11)$$

Therefore, the ground state wave function is given by $\Psi_0(\vartheta) = e^{iK\vartheta}/\sqrt{2\pi}$, with K an integer that is as close as possible to $\langle \Delta S_{||}^{\text{tot}} \rangle$. In this way we thus recover the fact that according to quantum mechanics the total number of spin-flips associated with the skyrmion texture must be an integer. More precisely, we have actually shown that the many-body wave function describing the skyrmion is an eigenstate of the operator $\hat{\mathbf{B}} \cdot \mathbf{S}^{\text{tot}}$ with eigenvalue $NS - K$. Note that physically this is equivalent to the way in which ‘diffusion’ of the overall phase of a Bose-Einstein condensate leads to the conservation of particle number [17,18].

Furthermore, the existence of this internal degree of freedom becomes especially important when we deal with more than one skyrmion in the condensate. In that case every skyrmion can have its own orientation and we expect the interaction between two skyrmions to have a Josephson-like contribution proportional to $\cos(\vartheta_1 - \vartheta_2)$. As a result the phase diagram of a gas of skyrmions can become extremely rich and contain both quantum as well as classical, i.e., nonzero temperature, phase transitions [16]. In this context, it is interesting to mention two important differences with the situation in the quantum Hall effect. First, the fact that the spin projection K of the skyrmion is an integer shows that these excitations have an integer spin and are therefore bosons [19]. In the quantum Hall case the skyrmions are fermions with half-integer spin, due to the presence of a topological term in the action $S[\zeta]$ for the spin texture [20]. Second, in a spinor Bose-Einstein condensate the skyrmions are not pinned by disorder and are in principle free to move. Both differences will clearly have important consequences for the many-body physics of a skyrmion gas.

Focussing again on the single skyrmion dynamics, we can make the last remark more quantitative by using the ansatz $\zeta(\mathbf{x}, t) = \zeta^{\text{sk}}(\mathbf{x} - \mathbf{u}(t))$ for the texture of a moving skyrmion, which is expected to be accurate for small velocities $\partial \mathbf{u}(t)/\partial t$. Considering for illustrative purposes also only the case of a maximally symmetric skyrmion near the center of the trap with $u/\xi \ll 1$, we find in a similar way as before that the action for the center-of-mass motion of the skyrmion becomes

$$S[\mathbf{u}] = \int dt \frac{1}{2} M \left(\frac{\partial \mathbf{u}(t)}{\partial t} \right)^2, \quad (12)$$

where the mass, which in the more general anisotropic case is of course a tensor, is now simply given by

$$M = \frac{m^2}{3} \int d\mathbf{x} \int d\mathbf{x}' \chi(\mathbf{x}, \mathbf{x}') \langle \mathbf{v}_s \rangle(\mathbf{x}) \cdot \langle \mathbf{v}_s \rangle(\mathbf{x}') \quad (13)$$

in terms of the superfluid velocity of the spinor condensate $\langle \mathbf{v}_s \rangle(\mathbf{x}) \equiv -i\hbar \zeta^{\text{sk}*}(\mathbf{x}) \nabla \zeta^{\text{sk}}(\mathbf{x})/m$. The skyrmions thus indeed behave in this respect as ordinary particles. It should be noted that in contrast to Eq. (9) there appears no term linear in $\partial \mathbf{u}(t)/\partial t$ in the action $S[\mathbf{u}]$. This is a result of the fact that we have performed all our calculations at zero temperature. In the presence of a normal component, we anticipate the appearance of such a linear term with an imaginary coefficient. This will lead to damping of the center-of-mass motion of the skyrmion.

I would like to thank Michiel Bijlsma, Gerard 't Hooft, David Olive, and Jan Smit for useful discussions. This work is supported by the Stichting Fundamenteel Onderzoek der Materie (FOM), which is financially supported by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO).

-
- [1] D.M. Stamper-Kurn, M.R. Andrews, A.P. Chikkatur, S. Inouye, H.-J. Miesner, J. Stenger, and W. Ketterle, Phys. Rev. Lett. **80**, 2027 (1998).
 - [2] H.-J. Miesner, D.M. Stamper-Kurn, J. Stenger, S. Inouye, A.P. Chikkatur, and W. Ketterle, Phys. Rev. Lett. **82**, 2228 (1999); D.M. Stamper-Kurn, H.-J. Miesner, A.P. Chikkatur, S. Inouye, J. Stenger, and W. Ketterle, Phys. Rev. Lett. **83**, 661 (1999)
 - [3] C.J. Myatt, A.E. Burt, R.W. Ghrist, E.A. Cornell, and C.E. Wieman, Phys. Rev. Lett. **78**, 586 (1997).
 - [4] M.R. Matthews, B.P. Anderson, P.C. Haljan, D.S. Hall, C.E. Wieman, and E.A. Cornell, Phys. Rev. Lett. **83**, 2498 (1999); M.R. Matthews, B.P. Anderson, P.C. Haljan, D.S. Hall, M.J. Holland, J.E. Williams, C.E. Wieman, and E.A. Cornell, Phys. Rev. Lett. **83**, 3358 (1999).
 - [5] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
 - [6] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
 - [7] C.K. Law, H. Pu, and N.P. Bigelow, Phys. Rev. Lett. **81**, 5257 (1998).
 - [8] T.-L. Ho and S.-K. Yip, cond-mat/9905339.
 - [9] S.-K. Yip, Phys. Rev. Lett. **83**, 4677 (1999).
 - [10] T.H.R. Skyrme, Proc. Roy. Soc. A **260**, 127 (1961); Nucl. Phys. **31**, 556 (1962).
 - [11] In principle, this is only true if the inter and intra scattering lengths of the two condensate components are equal, which is a good approximation for ^{87}Rb [4].
 - [12] N.D. Mermin, Rev. Mod. Phys. **51**, 591 (1979).
 - [13] G. 't Hooft, Nucl. Phys. B **79**, 276 (1974); A.M. Polyakov, JETP Lett. **20**, 194 (1974).
 - [14] H.T.C. Stoof, J. Stat. Phys. **87**, 1353 (1997) and references therein.
 - [15] S.L. Sondhi, A. Karlhede, S.A. Kivelson, and E.H. Rezayi, Phys. Rev. B **47**, 16419 (1993).
 - [16] R. Côté, A.H. MacDonald, L. Brey, H.A. Fertig, S.M. Girvin, and H.T.C. Stoof, Phys. Rev. Lett. **78**, 4825 (1997).
 - [17] M. Lewenstein and L. You, Phys. Rev. Lett. **77**, 3489 (1996).
 - [18] H.T.C. Stoof, J. Low Temp. Phys. **114**, 11 (1999).
 - [19] D. Finkelstein and J. Rubinstein, J. Math. Phys. **9**, 1762 (1968).
 - [20] F. Wilczek and A. Zee, Phys. Rev. Lett. **51**, 2250 (1983).